



DRUG FACTS TEXT (ABBREVIATED) TYPE SIZE/STYLE	
DRUG FACTS TITLE	N/A / N/A
HEADINGS	8 pt. / Helvetica Condensed Bold Obd.
SUBHEADS	6.0 pt. / Helvetica Condensed Bold
BODY TEXT	6.0 pt. / Helvetica Condensed
TRACKING #	HORIZONTAL SCALE: 85% LEADING: 6.5 pt.





NDC 0067-6286-28

**2 BOTTLES  
INSIDE**

**28 CAPSULES  
TOTAL**

# PREVACID®

**24 HR**

**PREVACID®**  
Lansoprazole delayed-release  
capsules 15 mg / acid reducer

**Actual Size**

ONE 14-DAY COURSE  
OF TREATMENT

Sodium Free

#### Drug Facts (continued)

##### Other information

- read the directions, warnings and package insert before use.

They contain important information.

• store at 20–25°C (68–77°F).

• keep product out of reach of heat and humidity.

• protect product from moisture.

**Inactive ingredients:** colloidal silicon dioxide, DSC, ref. No. 28, FD&C blue No. 1, FD&C green No. 3, FD&C red No. 40, gelatin, hydroxypropyl cellulose, low substituted hydroxypropyl cellulose, magnesium carbonate, methacrylic acid copolymer, polyethylene glycol, talc, titanium dioxide

##### Questions or comments?

Visit us at [www.prevacid24hr.com](http://www.prevacid24hr.com)

1-800-452-0051

##### KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

##### TAMPER-EVIDENT BOTTLE.

DO NOT USE IF NEVER FOIL SEAL IMPRINTED WITH "SESEALED" OR FOIL PROTECTION ON, BLACK BAND AROUND THE CENTER OF EACH CAPSULE IS MISSING OR BROKEN.

#### Drug Facts (continued)

##### Purpose

Acid Reducer

##### Uses

Lansoprazole 15 mg

##### Stop using and ask a doctor if:

• your heartburn continues or worsens.

• you need to take this product for more than 14 days.

• you need to take more than 1 course of treatment every 4 months.

• you get diarrhea.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a poison control center right away.

##### DirecTos

adults 18 years of age and older

This product is to be used once a day (every 24 hours), every day for 14 days.

• it may take 1 to 4 days to feel full relief, although some people get complete relief of symptoms within 24 hours.

##### 14-Day Course of Treatment

swallow 1 capsule with a glass of water before eating in the morning.

• take every day for 14 days.

• do not take more than 1 capsule a day.

• swallow whole. Do not crush or chew capsules.

##### Repeated 14-Day Courses (If needed)

• you may repeat a 14-day course every 4 months.

• do not take for more than 14 days or more often than every 4 months unless directed by a doctor.

• children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

##### Ask a doctor before use if you have:

• liver disease.

• bad heartburn over 3 months. This may be a sign of a more serious condition.

• heartburn with tightness, sweating or dizziness.

• chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or pain/tightness

frequent chest pain.

• frequent belching, particularly with heartburn.

• unexplained weight loss • nausea or vomiting • stomach pain

##### Ask a doctor or pharmacist before use if you are taking:

• warfarin (blood-thinning medicine)

• prescription artificial or anti-reject medicines

• digoxin (heart medicine)

• mequitazine (asthma medicine)

• tacrolimus or mycophenolate mofetil (immune system medicines)



DRUG FACTS TEXT	
DRUG FACTS TITLE	9.0 pt. / Helvetica Condensed Bold Oblique
HEADINGS	8.0 pt. / Helvetica Condensed Bold Oblique
SUBHEADS	6.0 pt. / Helvetica Condensed Bold
BODY TEXT	6.0 pt. / Helvetica Condensed
BULLETS	5.0 pt. / Helvetica Condensed
TRACKING 0	HORIZONTAL SCALE: 85% LEADING: 6.5 pt.
CHARACTERS/INCH: <39	
BARLINES, HAIRLINES: 1.5 pt., 0.5 pt.	


19.1

19.1

111.1 — 38.1 — 80.0 — 37.3 — 55.1 — 110.3 — 38.1 — 19.1



← →  $6\frac{1}{2}$



TAMPER-EVIDENT BOTTLE.  
DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED FOR YOUR PROTECTION"  
OR BLACK BAND AROUND THE CENTER OF EACH CAPSULE IS MISSING OR BROKEN.

**Drug Facts**

**Active ingredient (in each capsule)** Purpose  
Lansoprazole 15 mg ..... Acid reducer

**Uses**

- treats frequent heartburn (occurs 2 or more days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

**Warnings**

**Allergy alert:** Do not use if you are allergic to lansoprazole

**Do not use**

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

**Ask a doctor before use if you have**

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain      • frequent wheezing, particularly with heartburn
- unexplained weight loss      • nausea or vomiting      • stomach pain

**Ask a doctor or pharmacist before use if you are taking**

- warfarin (blood-thinning medicine)      • prescription antifungal or anti-yeast medicines
- digoxin (heart medicine)      • theophylline (asthma medicine)
- tacrolimus or mycophenolate mofetil (immune system medicines)
- atazanavir (medicine for HIV infection)      • methotrexate (arthritis medication)

**Stop use and ask a doctor if**

- your heartburn continues or worsens      • you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Visit us at [www.prevacid24hr.com](http://www.prevacid24hr.com)

**Drug Facts (continued)****Directions**

- adults 18 years of age and older
  - this product is to be used once a day (every 24 hours), every day for 14 days
  - it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

**14-Day Course of Treatment**

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

**Repeated 14-Day Courses (If needed)**

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor

children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

**Other information**

- read the directions, warnings and all information before use
- keep the package. It contains important information.
- store at 20-25°C (68-77°F)      • keep product out of high heat and humidity
- protect product from moisture

**Inactive ingredients** colloidal silicon dioxide, D&C red No. 28, FD&C blue No. 1, FD&C green No. 3, FD&C red No. 40, gelatin, hydroxypropyl cellulose, low substituted hydroxypropyl cellulose, magnesium carbonate, methacrylic acid copolymer, polyethylene glycol, polysorbate 80, starch, sucrose, sugar sphere, talc, titanium dioxide

**Questions or comments?** 1-800-452-0051



FPO

3 0067-6286-43 3

8<sup>15</sup>/<sub>16</sub>

**What you should know about PREVACID®24HR**

Please read all information before taking PREVACID®24HR.

Save for future reference.

**How PREVACID®24HR Treats Your Frequent Heartburn**

PREVACID®24HR stops acid production at the source – the pumps that release acid into the stomach. PREVACID®24HR is taken once a day (every 24 hours), every day for 14 days.

**What You Can Expect When Taking PREVACID®24HR**

Frequent heartburn can occur anytime during the 24-hour period (day or night). Take PREVACID®24HR in the morning before eating. PREVACID®24HR is clinically proven to treat frequent heartburn. Although some people get complete relief of symptoms within 24 hours, it may take 1 to 4 days for full effect. Make sure you take PREVACID®24HR every day for 14 days to treat your frequent heartburn.

**Who Should Take PREVACID®24HR**

Adults (18 years and older) with frequent heartburn – when you have heartburn 2 or more days a week.

**Who Should NOT Take PREVACID®24HR**

People who have one episode of heartburn a week or less, or who want immediate relief of heartburn.

**How to Take PREVACID®24HR****14-DAY Course of Treatment**

- Swallow 1 capsule with a glass of water before eating in the morning.
- Take every day for 14 days.
- Do not take more than 1 capsule a day.
- Swallow whole. Do not crush or chew capsules.
- Do not use for more than 14 days unless directed by your doctor.

**When to Take PREVACID®24HR Again**

You may repeat a 14-day course of therapy every 4 months.

**When to Talk To Your Doctor**

Do not take for more than 14 days or more often than every 4 months unless directed by a doctor.

**Warnings and When to Ask Your Doctor**

**Allergy alert:** Do not use if you are allergic to lansoprazole

**Do not use**

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

**Ask a doctor before use if you have**

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain      • frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

**Ask a doctor or pharmacist before use if you are taking**

- warfarin (blood-thinning medicine)
- prescription antifungal or anti-yeast medicines
- digoxin (heart medicine)
- theophylline (asthma medicine)
- tacrolimus or mycophenolate mofetil (immune system medicines)
- atazanavir (medicine for HIV infection)
- methotrexate (arthritis medication)

**Stop use and ask a doctor if**

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Tips for Managing Heartburn**

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.

- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.

**Clinical studies prove PREVACID®24HR effectively treats frequent heartburn**

In three clinical studies, PREVACID®24HR was shown to be significantly better than placebo in treating frequent heartburn.

**How PREVACID®24HR is Sold**

PREVACID®24HR is available in 14 capsule, 28 capsule and 42 capsule sizes. These sizes contain one, two and three 14-day courses of treatment, respectively. Do not use for more than 14 days in a row unless directed by your doctor. For the 28 count (two 14-day courses) and the 42 count (three 14-day courses), you may repeat a 14-day course every 4 months.

**For Questions or Comments About PREVACID®24HR**

Call 1-800-452-0051 or visit us at [www.prevacid24hr.com](http://www.prevacid24hr.com)

LOT:

EXP:

MADE IN:

DRUG FACTS (STANDARD) TEXT	
	TYPE SIZE/STYLE
DRUG FACTS TITLE	9.0 pt. / Helvetica Condensed Bold Oblique
HEADINGS	8.0 pt. / Helvetica Condensed Bold Oblique
SUBHEADS	6.0 pt. / Helvetica Condensed Bold
BODY TEXT	6.0 pt. / Helvetica Condensed
BULLETS	5.0 pt. / Helvetica Condensed
TRACKING: 0	HORIZONTAL SCALE: 95%
	LEADING: 6.5 pt.
CHARACTERS/INCH: <39	
BARLINES, HAIRLINES: 1.5 pt., 0.5 pt.	



NOVARTIS

Distributed by: Novartis Consumer Health, Inc., Parsippany, NJ 07054-0622 ©2014 141121-CB

PREVACID® is a registered trademark of Takeda Pharmaceuticals North America, Inc., and is used under license.



DRUG FACT FORMATTING: STANDARD		
DRUG FACTS TEXT	TYPE SIZE	TYPE STYLE
DRUG FACTS TITLE	9 pt.	Helvetica Cond Bold Obl
DRUG FACTS CONT.	8 pt.	Helvetica Cond Bold Obl
HEADINGS	8 pt.	Helvetica Cond Bold Obl
SUBHEADS	6 pt.	Helvetica Cond Bold
BODY TEXT	6 pt.	Helvetica Cond
TRACKING	0	
CONDENSATION	85%	
LEADING	6.5 pt.	
BARLINES, HAIRLINES	1.5, 0.5 pt.	
CHARACTERS/INCH	<39	
BULLETS	5 pt.	



---

---

**This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.**

---

/s/

---

VALERIE S PRATT

06/30/2015